

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/335255804>

# Consumption of aquatic macrophytes by the Red-gartered Coot *Fulica armillata* (Aves: Rallidae) in a coastal wetland of north central Chile

Article in *Gayana (Concepción)* · June 2019

DOI: 10.4067/S0717-65382019000100068

CITATIONS

5

READS

105

4 authors, including:



**Carlos Velásquez**

16 PUBLICATIONS 99 CITATIONS

[SEE PROFILE](#)



**Eduardo Jaramillo**

Universidad Austral de Chile

127 PUBLICATIONS 4,228 CITATIONS

[SEE PROFILE](#)



**Patricio A. Camus**

Universidad Católica de la Santísima Concepción

104 PUBLICATIONS 2,265 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



CYCLO The seismic cycle along subduction zones [View project](#)

## Consumption of aquatic macrophytes by the Red-gartered Coot *Fulica armillata* (Birds: Rallidae) in a coastal wetland of north central Chile

### Consumo de macrófitas acuáticas por la Tagua común *Fulica armillata* (Aves: Rallidae) en un humedal costero del centro norte de Chile

CARLOS VELÁSQUEZ<sup>1,2\*</sup>, EDUARDO JARAMILLO<sup>3</sup>, PATRICIO A. CAMUS<sup>4</sup> & CRISTINA SAN MARTÍN<sup>3</sup>

<sup>1</sup>Programa de Magíster en Ciencias mención Recursos Hídricos, Escuela de Graduados, Facultad de Ciencias, Universidad Austral de Chile, Casilla 567, Valdivia, Chile.

<sup>2</sup>Instituto de Fomento Pesquero, Dr. Marín 340, Coquimbo, Chile.

<sup>3</sup>Instituto de Ciencias de la Tierra, Facultad de Ciencias, Universidad Austral de Chile, Casilla 567, Valdivia, Chile.

<sup>4</sup>Departamento de Ecología, Facultad de Ciencias y Centro de Investigación en Biodiversidad y Ambientes Sustentables CIBAS, Universidad Católica de la Santísima Concepción, CP 4090541, Concepción, Chile.

\*cfvelasque@gmail.com

#### ABSTRACT

The Red-gartered Coot *Fulica armillata*, is a common herbivorous water bird in coastal wetlands of the southern Neotropical region. Microhistological analyses of feces collected at the coastal wetland of Punta Teatinos (north central Chile; ~29°S), show that this coot feeds predominantly on *Stuckenia pectinata*, the most abundant submerged macrophyte at this area. The foraging behavior of *F. armillata* is discussed in regard with its ecological role as a primary consumer on coastal wetlands.

#### RESUMEN

La Tagua común *Fulica armillata*, es un ave acuática herbívora común en los humedales costeros del extremo sur de la región del Neotrópico. Análisis microhistológicos de fecas recolectadas en el humedal de Punta Teatinos (centro norte de Chile; ~29°S) indican que esta tagua consume primariamente a *Stuckenia pectinata*, la macrófita sumergida más abundante en esta área. Se discute el comportamiento de forrajeo de *F. armillata* con respecto a su rol ecológico como consumidor primario en humedales costeros.

---

The Red-gartered Coot *Fulica armillata* (Vieillot, 1817) (Aves: Rallidae) is one of the six species of this genus occurring in limnetic wetlands of the southern Neotropical region, including Chile, Argentina, Uruguay and southeastern Brazil (Couve *et al.* 2016). In Chile, this coot has a wide latitudinal distribution: from the Río Copiapó in the north (~27°S) to Isla Grande de Tierra del Fuego and Isla Navarino in the south (~54°S) (Medrano *et al.* 2018). This species has been described as a primarily herbivorous waterbird, although its diet also includes invertebrates although in lower proportion than plants (Ruíz 1993; García *et al.* 2008).

The abundance of *F. armillata* in the southern Neotropical region has been estimated in ~1,000,000 birds (Waterbird Population Estimates 2017). Thus, and due to its herbivorous feeding habits, this bird may potentially play an important role both as controller of the biomass of aquatic macrophytes and as seed disperser (Ruíz 1993; Bortolus *et al.* 1998; Charalambidou & Santamaría 2005),

as shown for other coots such as *Fulica cristata* (Gmelin, 1789) (Stewart & Bally 1985), *Fulica atra* (Linnaeus, 1758) (Perrow *et al.* 1997) and *Fulica americana* (Gmelin, 1789) (Esler 1989). Despite the important ecological role of coots, the natural history of *F. armillata* in Chile is scarcely known. Cody (1970) and Riveros *et al.* (1981) studied the spatial distribution of this bird in wetlands of central Chile (c. 30–38°S), Silva *et al.* (2011) described its reproductive biology in Laguna Santa Elena (central Chile; ~36°S), while Kennedy (1977) and Ruíz (1993) studied ecological and trophic aspects at the wetland of Río Cruces (southern Chile; ~39°S). However, no studies have been carried out on populations of *F. armillata* along the semiarid coastal region of northern Chile, where coastal wetlands host a different suite of waterbirds compared to their southern counterparts (Sielfeld *et al.* 2012). An important wetland at that region is Punta Teatinos (PT hereafter), located north of the conurbation of Coquimbo and La Serena in north central Chile (~29°S; Fig. 1a, b), where field observations carried

out during 2016–17 showed that *F. armillata* is one of the most abundant resident herbivorous waterbirds. Therefore, this study was aimed to provide a first assessment of the use of plant food resources by *F. armillata* at the PT wetland, based on a microhistological analysis of its feces and a comparison with published results.

The PT wetland is a coastal lagoon of ~69 ha with freshwater inflows from the ravine El Romeral and salty water influence from the nearby sand beach surf zone during extreme high tides (Jorge *et al.* 1998). *Stuckenia pectinata* (L.) Börner is the most abundant submerged macrophyte at this lagoon (C. Velásquez, obs. Pers.). The western side of the wetland (see Fig. 1c) is bordered by the upper shore sandy beach level, which *F. armillata* used as dropping area. Thus, during May 2016, fresh feces of the Red-gartered Coot ( $n = 30$ ) were randomly collected along the western sandy bank of PT, and then stored in sealed plastic bags with

alcohol (70 %) to preserve them. Additionally, with the aim of maximizing the likelihood that collected feces belonged to different coot individuals, the sampling was carried out through the entire border of the lagoon, at places used by red-gartered coots as resting or passing spots.

In order to identify the species of plants in the feces, a series of microhistological techniques (after Johnson *et al.* 1983) to analyze food habits of herbivores were used to (i) prepare samples of epidermal cells collected from the macrophytes of the PT wetland, (ii) compare those samples with the plant epidermal cells found intact in coot feces, and (iii) quantify the examined material. The comparisons were carried out with an optic microscope (10x magnification) connected to the software Micrometrics Premium to obtain high resolution digital pictures (ACCU-SCOPE camera) (see Velásquez *et al.* 2018).




FIGURE 1. Geographic location of the wetland of Punta Teatinos, in north central Chile (a) and at the northern side of Bahía de Coquimbo (b). A general picture of the wetland is also included (c); the red arrows indicate the western side of the wetland where the feces of coots were collected. / Localización geográfica del humedal de Punta Teatinos, en el centro norte de Chile (a) y en el costado norte de Bahía de Coquimbo (b). Se incluye una foto general del área de estudio (c); las flechas rojas indican el lado oeste del humedal donde se recolectaron las fecas de las taguas.

The results were first expressed as the percent occurrence frequency of each plant species in the sample of 30 feces (replicates). However, this measure cannot clearly reflect the intensity of consumption of food items, considering that the presence of different plants in each feces is recorded only once, regardless of the amount of biomass ingested of each plant. Therefore, an exhaustive subsampling was conducted in order to obtain an estimate of the incidence of each plant species within the feces. To that purpose, two portions from each feces (herein referred to as subsamples) were randomly selected, and each subsample was uniformly spread out in a Neubauer counting chamber, where 10 optic fields of 1 mm<sup>2</sup> were also randomly selected for observation, thus examining a total of 600 fields (*i.e.* 10 fields x 2 chambers x 30 feces). The presence of each plant species in each set of 10 fields per chamber was recorded as a percent frequency, and the two frequency values recorded for each feces were then averaged. Finally, the incidence of each plant species was expressed as the grand average ( $\pm$  SE) of the 30 average values obtained from the feces collected in the PT wetland.

The analysis of feces showed that the only conspicuous macrophytes consumed by *F. armillata* were *Cotula coronopifolia* L., *Sarcocornia fruticosa* (L.) Scott, and *S. pectinata* which was by far the dominant food item (Fig. 2). Remarkably, *S. pectinata* was present not only at all feces, with an occurrence frequency of 100.0 %, but also at each of the 600 optic fields examined, with an incidence of  $100.0 \pm 0.0$  % (Table 1) suggesting a heavy consumption by this species. In contrast, the occurrence frequency of *C. coronopifolia* and *S. fruticosa* in the 30 feces was 66.7 % and 36.7 %, respectively, but their incidence within the feces was much lower, reaching only  $13.9 \pm 0.6$  % and  $15.3 \pm 1.0$  %, respectively (Table 1). In addition, these results were consistent with the different life habits of the three macrophytes (see Ramírez & San Martín 2006), indicating that coots fed predominantly on submerged (*S. pectinata*)

rather than marshland (*C. coronopifolia* and *S. fruticosa*) plants (Table 1).

Albeit the above results do not allow a proper evaluation of feeding preferences (*e.g.* Jacksic *et al.* 1979), prior antecedents suggest that the high dietary importance of *S. pectinata* would not be due to its selective consumption by coots, but rather to its submerged habit and its high abundance at the PT wetland. Unlike other Chilean coots such as *Fulica leucoptera* (Vieillot, 1817) and *Fulica rufifrons* (Philippi & Landbeck, 1861), *F. armillata* is known to display its foraging activities mainly at the central, deeper portion of lagoons (Cody 1970; Riveros *et al.* 1981). On the other hand, in shallower, non-lagoon wetlands such as the large wetland of Río Cruces (southern Chile), both *F. armillata* and *F. leucoptera* (see Table 1) feed primarily (~85 %) on the submerged and most abundant macrophytes (particularly *Egeria densa* Planch; Ramírez *et al.* 1991; Ruíz 1993), a pattern shown also by other common water birds such as the Black-necked Swan *Cygnus melancoryphus* (Molina, 1782) (Table 1; Corti & Schlatter 2002; Norambuena & Bozinovic 2009). Thus, *F. armillata* would behave as an opportunistic consumer, although future studies should clarify whether the submerged habit itself is a factor influencing the feeding choice of this species.

On a general level, the similar foraging patterns of the commonest herbivorous birds across wetlands of Chile and the southern Neotropical region (Couve *et al.* 2016) might play an important role in the succession of macrophyte assemblages. Submerged macrophytes such as *S. pectinata*, *E. densa* and *Ruppia maritima* L. are dominant in coastal wetlands of the southern Neotropic, and the large amount of plant necromass derived from bird foraging may determine the zonation and successional patterns of the vegetation, or even retard the successional process (Ruíz 1993; Bortolus *et al.* 1998; Corti & Schlatter 2002).



FIGURE 2. Pictures of the aquatic macrophytes consumed by *Fulica armillata* in the study area: *Stuckenia pectinata* (a), *Cotula coronopifolia* (b) and *Sarcocornia fruticosa* (c). / Fotografías de las macrófitas acuáticas consumidas por *Fulica armillata* en el área de estudio: *Stuckenia pectinata* (a), *Cotula coronopifolia* (b) and *Sarcocornia fruticosa* (c).

TABLE 1. Frequency of occurrence of the food items of herbivorous waterbirds in coastal wetlands of Chile (*Fulica armillata*, *Fulica leucoptera* and *Cygnus melancoryphus*). The macrophytes are classified by their life habits (see Ramírez & San Martín 2006). Food items were determined throughout analyses of feces (\*) and analyses of stomach contents (\*\*). / Frecuencia de ocurrencia del ítem alimentario de aves acuáticas herbívoras en humedales costeros de Chile (*Fulica armillata*, *Fulica leucoptera* y *Cygnus melancoryphus*). Las macrófitas se clasifican según sus hábitos de vida (ver Ramírez & San Martín 2006). Los ítems alimentarios fueron determinados a través del análisis de fecas (\*) y análisis de contenido estomacal (\*\*).

FOOD ITEMS (%)	THIS STUDY*	Ruíz (1993)**		Corti & Schlatter (2002)*		Norambuena & Bozinovic (2009)*
	<i>F. armillata</i> P. Teatinos	<i>F. armillata</i> Río Cruces	<i>F. leucoptera</i> Río Cruces	<i>C. melancoryphus</i> Río Cruces    Chihuahua		<i>C. melancoryphus</i> Lago Budi
	n = 30	n = 18	n = 7	n = 13	n = 8	n = 10
<b>Submerged</b>						
<i>Stuckenia pectinata</i>	100.0					
<i>Stuckenia striata</i>						78.0
<i>Potamogeton pusillus</i>		2.2	1.0	7.6	1.3	
<i>Egeria densa</i>		85.3	84.8	91.9	71.7	
<b>Free floating</b>						
<i>Limnobium laevigatum</i>				< 1.0	24.7	
<b>Floating</b>						
<i>Ludwigia peploides</i>					< 1.0	
<i>Hydrocotyle ranunculoides</i>					2.1	
<b>Marshland</b>						
<i>Puccinellia glaucescens</i>						20.5
<i>Cotula coronopifolia</i>	66.7					< 1.0
<i>Sarcocornia fruticosa</i>	36.7					< 1.0

Thus, the close connection between submerged macrophytes and herbivorous waterbirds such as *F. armillata* would make wetlands highly sensitive to bottom-up forcing. For instance, industrial activities triggered sudden environmental changes at the Río Cruces wetland in 2004, leading to a radical decline in the spatial cover of *E. densa* and the population abundance of *F. armillata* and others birds (e.g. Jaramillo *et al.* 2007; Lagos *et al.* 2008). Only eight years later, macrophytes begun to recover and a concomitant increase in the abundance of coots was observed (Jaramillo *et al.* 2018a, 2018b), although the mechanisms underlying their decline and subsequent recovery remain not fully understood.

Even though *F. armillata* is not considered an endangered species (Bird Life International 2017), a better knowledge

of its trophic ecology may provide crucial information for the management and restoration of coastal wetlands over the southern Neotropical region.

## ACKNOWLEDGEMENTS

We thank to Felipe Dreves (<http://www.lluviafilms.com>), Wolfgang Stotz (Universidad Católica del Norte, Chile) and Emilio Acuña (Ecosistema Ltda.) for logistical support. CV thanks the support of Escuela de Graduados, Facultad de Ciencias (UACH), Vicerrectoría de Investigación & Creación Artística (Proyecto: I-2015-10; UACH) and Beca de Asistencia Académica de la Dirección de Postgrado (2015-2016; UACH).

REFERENCES

- WATERBIRD POPULATION ESTIMATES. 2017. Species factsheet: *Fulica armillata*. URL: <http://www.wpe.wetlands.org>. Accessed: December 5, 2018.
- BIRDLIFE INTERNATIONAL. 2017. Species factsheet: *Fulica armillata*. URL: <http://www.birdlife.org>. Accessed: October 15, 2017.
- BORTOLUS, A., IRIBARNE, O.O., MARTÍNEZ, M.M. 1998. Relationship between waterfowl and the seagrass *Ruppia maritima* in a southwestern Atlantic coastal lagoon. *Estuaries* 21: 170-717.
- CHARALAMBIDOU, I., SANTAMARÍA, L. 2005. Field evidence for the potential of waterbirds as dispersers of aquatic organisms. *Wetlands* 25(2): 252-258.
- CODY, M. 1970. Chilean bird distribution. *Ecology* 59:455-464.
- CORTI, P., SCHLATTER, R. 2002. Feeding ecology of Black-necked swan *Cygnus melancoryphus* in two wetlands of Southern Chile. *Studies on Neotropical Fauna and Environment* 37: 9-14.
- COUVE, E., VIDAL, C.F., RUÍZ, J. 2016. Aves de Chile, sus Islas Oceánicas y Península Antártica. FS Editorial, Punta Arenas. 551 pp.
- ESLER, D. 1989. An assessment of American coot herbivory of *Hydrilla*. *Journal of Wildlife Management* 53: 1147-1149.
- GARCÍA, G.O., FAVERO, M., MARIANO-JELICICH, R. 2008. Red-gartered Coot *Fulica armillata* feeding on the grapsid crab *Cyrtograpsus angulatus*: advantages and disadvantages of an unusual food resource. *Ibis* 150: 110-114.
- JAKSIC, J.M. 1979. Técnicas estadísticas simples para evaluar selectividad dietaria en Strigiformes. *Medio Ambiente* 4: 114-118.
- JARAMILLO, E., SCHLATTER, R., CONTRERAS, H., DUARTE, C., LAGOS, N., PAREDES, E., ULLOA, J., VALENZUELA, G., PERUZZO, B., SILVA, R. 2007. Emigration and mortality of black-necked swans (*Cygnus melancoryphus*) and disappearance of the macrophyte *Egeria densa* in a Ramsar wetland site of southern Chile. *Ambio* 36: 607-610.
- JARAMILLO, E., LAGOS, N.A., LABRA, F.A., PAREDES, E., ACUÑA, E.O., MELNICK, D., MANZANO, M., VELÁSQUEZ, C., DUARTE, C. 2018a. Recovery of Black-necked Swans, macrophytes and water quality in a Ramsar wetland of southern Chile: assessing resilience following sudden anthropogenic disturbances. *Science of the Total Environment* 628: 291-301.
- JARAMILLO, E., DUARTE, C., LABRA, F.A., LAGOS, N.A., PERUZZO, B., VELÁSQUEZ, C., MANZANO, M., MELNICK, D., SILVA, R. 2018b. Resilience of an aquatic macrophyte to an anthropogenically induced environmental stressor in a Ramsar wetland of southern Chile. *AMBIO* DOI: 10.1007/s13280-018-1071-6.
- JOHNSON, M.K., WOFFORD, H., PEARSON, H.A. 1983. Microhistological techniques for food habits analyses. Forest Service, New Orleans. 40 pp.
- JORGE, R., TABILO-VALDIVIESO, E., MONCADA, V. 1998. Avifauna de la Laguna de Punta Teatinos y ecosistemas adyacentes, Bahía de Coquimbo, Chile. *Boletín Chileno de Ornitología* 5: 2-9.
- KENNEDY, M.E. 1977. Requisitos para el hábitat de reproducción de la tagua común (*Fulica armillata*) y recomendaciones técnicas de manejo para aumentar esta especie. *Medio Ambiente* 2: 107-116.
- LAGOS, N.A., PAOLINI, P., JARAMILLO, E., LOVINGREEN, C., DUARTE, C., CONTRERAS, H. 2008. Environmental processes, water quality degradation, and decline of waterbird populations in the Río Cruces wetland, Chile. *Wetlands* 28: 938-950.
- MEDRANO, F., BARROS, R., NORAMBUENA, H.V., MATUS, R., SCHMITT, F. 2018. Atlas de las aves nidificantes de Chile. Red de Observadores de Aves y Vida Silvestre de Chile, Santiago. 658 pp.
- NORAMBUENA, C.M., BOZINOVIC, F. 2009. Health and nutritional status of a perturbed Black-necked swan (*Cygnus melancoryphus*) population: diet quality. *Journal of Zoo and Wildlife Medicine* 40: 607-616.
- PERROW, M.R., SCHUTTEN, J.H., HOWES, J.R., HOLZER, T., MADGWICK, F.J., JOWITT, A.J. 1997. Interactions between coot (*Fulica atra*) and submerged macrophytes: the role of birds in the restoration process. *Hydrobiologia* 342/343: 241-255.
- RAMÍREZ, C., SAN MARTÍN, C., MEDINA, R., CONTRERAS, D. 1991. Estudio de la flora hidrófila del Santuario de la Naturaleza "Río Cruces" (Valdivia, Chile). *Gayana Botanica* 48: 67-80.
- RAMÍREZ, C., SAN MARTÍN, C. 2006. Diversidad de macrófitos chilenos. En: Vila, I., Veloso, A., Schlatter, R., Ramírez, C. (Eds.). *Macrófitas y vertebrados de los sistemas límnicos de Chile*. Editorial Universitaria, Santiago, Chile. pp. 21-60.
- RIVEROS G., SEREY, I., DROUILLY, P. 1981. Estructura y diversidad de la comunidad de aves acuáticas de la laguna El Peral, Chile Central. *Anales del Museo de Historia Natural* 14: 189-196.
- RUIZ, J. 1993. Estudio ecológico en tres especies de taguas residentes en el Santuario de la Naturaleza del río Cruces. Tesis. Universidad Austral de Chile, Valdivia, Chile.
- SILVA, C., BARRIENTOS, C., FIGUEROA, R.A., MARTÍN, N., CONTRERAS, A., ARDILES, K., MORENO, L., GONZÁLEZ-ACUÑA, D. 2011. Biología reproductiva de la tagua común (*Fulica armillata*) y la tagua de frente roja (*Fulica rufifrons*) en un área agroforestal del centro-sur de Chile. *Gayana* 75: 161-169.
- SIELFELD, W., FUENTES, R., PEREDO, R., MALINARICH, V., OLIVARES, F. 2012. Humedales costeros del norte de Chile. En: Humedales costeros de Chile. Aportes científicos a su gestión sustentable (Eds. Fariña J.M., Camaño, A.), pp. 147-213. Ediciones Universidad Católica de Chile, Santiago.
- STEWART, B.A., BALLY, R. 1985. The ecological role of the Red-knobbed Coot *Fulica cristata* Gmelin at the Bot River Estuary South Africa: a preliminary investigation. *Transactions of the Royal Society of South Africa* 45: 419-426.
- VELÁSQUEZ, C., SAN MARTÍN, C., JARAMILLO, E., CAMUS P.A. 2018. Catálogo microhistológico de macrófitas acuáticas de dos humedales costeros de Chile: Una herramienta para estudios tróficos en aves acuáticas herbívoras. *Revista Chilena de Ornitología* 24: 79-84.

Recibido: 05.12.2018  
Aceptado: 08.03.2019